



Perry Local Schools Immunization Requirements

Vaccine	
DTap/DTP/DT/Td/Tdap Diphtheria, Tetanus, Pertussis	<p>Kindergarten – Grade 12</p> <ul style="list-style-type: none"> • Four (4) or more of DTap or DT, or any combination. If fourth dose was given before 4th birthday, a fifth dose is required. <p>Grades 7 – 10</p> <ul style="list-style-type: none"> • 1 dose of Tdap or Td must be administered prior to school entry. <ul style="list-style-type: none"> ○ Persons 11 – 18 years who have not received Tdap should receive a dose. ○ Tdap can be administered regardless of the interval since the last tetanus and diphtheria toxoid-containing vaccine.
Polio	<p>Kindergarten – Grade 3</p> <ul style="list-style-type: none"> • Three (3) or more doses of IPV; the FINAL dose must be administered on or after the 4th birthday regardless of the number of previous doses. If a combination of OPV and IPV was received, 4 doses of either vaccine are required. <p>Grades 4 – 12</p> <ul style="list-style-type: none"> • Three (3) or more doses of IPV or OPV. If the third dose of either series was received prior to the fourth birthday, a fourth dose is required; If a combination of OPV and IPV was received, four (4) doses of either vaccine are required.
MMR Measles, Mumps, Rubella	<p>Kindergarten – Grade 12</p> <ul style="list-style-type: none"> • 2 doses of MMR • Dose 1 must be administered on or after the first birthday. • Dose 2 must be administered at least 28 days after Dose 1.
Hib Haemophilus Influenzae Type B	<p>REQUIREMENT FOR PRESCHOOL ONLY</p> <ul style="list-style-type: none"> • 3 or 4 doses depending on the vaccine type and the age when the child began the first dose and the last dose is after 12 months OR • 1 dose is given on or after 15 months of age.
Hep B Hepatitis B	<p>Kindergarten – Grade 12</p> <ul style="list-style-type: none"> • 3 doses of Hep B. The second dose must be administered at least 28 days after the first. • The third dose should be given at least 16 weeks after the first dose and at least 8 weeks after the second dose. • The last dose in the series (3rd or 4th dose) MAY NOT be administered before age 24 weeks.
Varicella Chickenpox	<p>Kindergarten – Grade 3</p> <ul style="list-style-type: none"> • 2 doses of varicella MUST be administered prior to school entry. <p>Grades 4 – 7</p> <ul style="list-style-type: none"> • 1 dose of varicella must be administered on or after first birthday.

Additional vaccines which your primary care giver may recommend are:

- Meningococcal (meningitis) vaccine – all 11 -12 year old should be vaccinated and receive a booster at approximately age 16.
- Hepatitis A vaccine – first dose at 12 – 23 months of age. Additional vaccines age 2 – 18 years.
- Human Papillomavirus Vaccine – recommended for preteen girls AND boys at age 11 – 12 years.
- Influenza vaccine yearly.